

Resolved Photons and BFKL-type Signatures in Deep Inelastic Scattering^a

H. Jung

University of Lund, Department of Physics, Sweden

The concept of resolved virtual photons in addition to direct deep inelastic ep scattering is used to simulate the $2+1$ jet - rate and the forward jet cross section, which cannot be described by direct LO/NLO processes. With standard DGLAP evolution of the parton densities of the virtual photon and the proton the HERA data can be described, provided a sufficiently large scale for the hard scattering matrix element is used.

1 Introduction

At HERA the measurement of the inclusive forward jet production^{1,2} in the proton direction in deep inelastic events at low x and large Q^2 is believed to be one of the most sensitive measurements for BFKL dynamics. The measured cross section³ cannot be explained by standard deep inelastic scattering models and even next - to - leading - order calculations (NLO, order α_s^2) predict a too small cross section⁴.

The measured⁵ fractional $2+1$ jet rate $R_{2+1} = \frac{\sigma^{2+1}}{\sigma^{tot}}$, where two high p_T jets come from the hard QCD scattering and the "1" stands for the remnant jet, seems to be a factor of ~ 2 larger than predictions from standard deep inelastic scattering (DIS) models.

However both measurements are rather well described by the Color - Dipole - Model (CDM) in its implementation in ARIADNE⁶.

In this article we study to what extent the contribution of virtual resolved photons in addition to the standard DIS matrix element calculations (labeled as *direct* contributions) can reproduce the data. The studies are performed with the RAPGAP⁷ Monte Carlo event generator which for the direct processes agrees rather well with the LEPTO⁸ Monte Carlo program.

2 Resolved Photons in DIS

Resolved photon processes play an important role in photo-production, when high p_T jets are observed. Any internal structure of the proton as well as of the photon can be resolved as long as the scale μ^2 of the hard subprocess, which is

^aTo be published in Proceedings of the *Madrid low x Workshop*, Miraflores de la Sierra, 1997

of the order of p_T^2 , is larger than the inverse radius of the proton $1/R_p^2 \sim \Lambda_{QCD}^2$ and the photon $1/R_\gamma^2 \sim Q^2$.

Resolved photon processes in DIS are calculated here using the Equivalent Photon Approximation, giving the flux of virtual transverse polarized photons^{7,9} together with the on shell matrix elements for the partonic subprocesses. The structure of the virtual photon is given by parametrisations of the parton densities^{10,11,12} $x_\gamma f_\gamma(x_\gamma, \mu^2, Q^2)$, now depending on the scale μ^2 and the photon virtuality Q^2 . Direct deep inelastic scattering and resolved virtual photon processes are simulated using the RAPGAP Monte Carlo event generator⁷. The hard subprocesses for resolved photons implemented in RAPGAP are: $gg \rightarrow q\bar{q}$, $gg \rightarrow gg$, $qg \rightarrow qg$, $q\bar{q} \rightarrow gg$, $q\bar{q} \rightarrow q\bar{q}$ and $qq \rightarrow qq$.

In DIS, resolved photon processes^{13,10} can play a role when the scale μ^2 of the hard subprocess is larger than the virtuality of the photon Q^2 : $\mu^2 > Q^2$. A typical NLO diagram like $\gamma^* g \rightarrow q\bar{q}g$ occurs also in the resolved photon picture, via the subprocess $\bar{q}g \rightarrow \bar{q}g$ with the photon splitting $\gamma^* \rightarrow q\bar{q}$. When including resolved photon processes in a NLO calculation special care has to be taken to avoid double counting¹⁴. In leading order there is no double counting when adding resolved photons to direct processes since they have different final states (three parton and two parton).

2.1 Scales, α_s and Parton Distribution Functions

In leading order α_s processes the renormalisation scale μ_R and factorization scale μ_F are not well defined, and any choice of $\mu_R^2 = \mu_F^2 = Q^2$, p_T^2 , $Q^2 + p_T^2$, and multiples thereof, is reasonable. There are essentially two competing effects: a larger scale gives smaller $\alpha_s(\mu^2)$, but a larger parton density $xf(x, \mu^2)$ at fixed x . The net effect depends on the details of the interaction.

The effect of different choices of the scale $\mu^2 = Q^2 + 4 \cdot p_T^2$, $\mu^2 = Q^2 + p_T^2$, $\mu^2 = 4 \cdot p_T^2$, or $\mu^2 = p_T^2$ is discussed in the following.

We use the GRV 94 HO (DIS) parametrisation of the proton parton densities^{15,16} together with the SaSgam 2D¹¹ parametrisation for the partons in the virtual photon. Other choices like the GRV LO parton density of the real photon together with the virtual photon suppression factor of Drees - Godbole¹² give similar results.

3 Forward Jets

We study the HERA data on high p_T “forward” jet production at low x and large Q^2 in the context of resolved virtual photons. The cuts are taken from the preliminary H1 analysis³: $y > 0.1$, $E'_e > 11$ GeV, $160^\circ < \theta_e < 173^\circ$,




Figure 1: The forward jet cross section as a function of x using the cuts of the H1 collaboration specified in the text. The data points are preliminary H1 data³. In 1a.) the dashed line is the RAPGAP prediction for direct DIS processes and the solid line is the RAPGAP prediction for the sum of direct and resolved contributions (using $\mu^2 = Q^2 + 4p_T^2$, GRV HO for the proton and SaSgam 2 DIS for the virtual photon). In 1b.) the solid line is the RAPGAP prediction for the sum of direct and resolved contributions with $\mu^2 = Q^2 + p_T^2$, the dashed line with $\mu^2 = 4p_T^2$ and the dotted line with $\mu^2 = p_T^2$. The difference between solid and dashed lines is hardly visible.

$7^\circ < \theta_{jet} < 20^\circ$, $E_{jet} > 28.7$ GeV, and $p_T^{jet} > 3.5$ GeV in the laboratory frame. The observed forward jet is required to have sufficiently large E_{jet} and p_T^{jet} in order to rely on well measured parton distributions in the proton. A p_T of the jet is required to be of the same order as Q^2 ($0.5 < p_T^2/Q^2 < 2$) to suppress the contribution from direct LO processes which are governed by DGLAP evolution. The event topology is a small x deep inelastic event with an energetic high p_T jet in the forward (proton) direction. In Fig. 1a.) preliminary data of the H1 collaboration³ are shown together with the RAPGAP⁷ Monte Carlo prediction for *standard* DIS processes (*direct*) and the sum of direct and resolved processes, using the scale $\mu^2 = Q^2 + 4p_T^2$. We note that the dominant contribution of resolved photon processes to the forward jet cross section comes from $qg \rightarrow qg$.

With a scale $\mu^2 = Q^2 + 4p_T^2$ (with p_T defined in the parton parton cms) a perfect description of the data (Fig. 1a.) is achieved. The effect of different choices of the scale μ^2 is shown in Fig. 1b.). We observe no difference using

$\mu^2 = Q^2 + p_T^2$ or $\mu^2 = 4p_T^2$. In general a less hard scale results in a smaller cross section and worse agreement with the data at small x values. Using $\mu^2 = p_T^2$ the predicted cross section falls well below the measurement. This is because the condition $\mu^2 = p_T^2 > Q^2$ together with $0.5 < Q^2/p_T^2 < 2$ leaves only little space for resolved photon contributions. However with the concept of resolved virtual photons we obtain a improved description of the data for all choices of the scale μ^2 .

4 2 + 1 jet rates in DIS

Dijets (2 + 1 - jets) are searched for with a cone jet algorithm by requiring a $p_T^{jets} > 5$ GeV in the hadronic center of mass frame, $Q^2 > 5$ GeV², $y > 0.05$, $E' > 11$ GeV and $156^\circ < \theta_e < 173^\circ$ according to the preliminary H1 - analysis⁵. Since in that analysis $p_T^2 > Q^2$ in the region $Q^2 < 25$ GeV², resolved photon interactions can be expected to play a significant role. It has been observed⁵, that *direct* LO Monte Carlos like RAPGAP *direct* and LEPTO as well as NLO calculations are unable to describe the data especially in the low Q^2 region of $Q^2 < 30$ GeV². However ARIADNE agrees with the data remarkably well. In Fig. 2a., and b.) the H1 measurement⁵ of R_{2+1} as a function of Q^2 and x is shown together with the *direct* contribution and the sum of *direct* and resolved photons obtained with RAPGAP. The parameter setting is the same as for the forward jet analysis. We find remarkable agreement (using $\mu^2 = Q^2 + 4p_T^2$) between the data and the Monte Carlo prediction showing that especially in the region of phase space where $Q^2 < p_T^2$, resolved photons play an important role. Note that here also the dominant contribution from resolved photon processes comes from $qg \rightarrow qg$.

We have checked R_{2+1} as a function of Q^2 and x also for different scales $\mu^2 = Q^2 + p_T^2$, $\mu^2 = 4p_T^2$ and $\mu^2 = p_T^2$ used for the resolved photon contribution. The contribution of resolved photons to R_{2+1} is larger for the largest scale $\mu^2 = Q^2 + 4p_T^2$, since there is a larger phase space for DGLAP evolution in the photon. The results are shown in Fig. 2c., and d.) Again we observe the same trend, that including resolved photons in DIS, the description of the measurements is considerably improved for all different choices of the scales μ^2 .

5 Discussion

A scale dependence is expected in all leading order calculations, but it becomes more important when the concept of resolved photons is applied to DIS: in order to resolve any internal structure of the virtual photon, the scale involved




Figure 2: The 2+1 jet ratio R_{2+1} as a function of Q^2 and x . The data points are preliminary H1 data⁵. In 2a.) and b.) the dotted line is the RAPGAP prediction for the direct contribution, the dashed line for resolved photon contribution and the solid line is the prediction for the sum of direct and resolved contributions (using $\mu^2 = Q^2 + 4p_T^2$, GRV HO for the proton and SaSgam 2 DIS for the virtual photon). In 2c.) and d.) the solid line is the RAPGAP prediction for the sum of direct and resolved contributions with $\mu^2 = Q^2 + p_T^2$, the dashed line with $\mu^2 = 4p_T^2$ and the dotted line with $\mu^2 = p_T^2$.

has to be larger than the virtuality of the photon Q^2 : $\mu^2 > Q^2$. Using a large scale like $\mu^2 = Q^2 + 4p_T^2$ results in a larger magnitude of the parton density, both on the photon and proton side, at fixed values of fractional parton momenta ξ_i , due to the larger phase space for DGLAP evolution. This effect is compensated to some extent by the smaller value of α_s at the same scale.

In case of the forward jet cross section the stronger scale dependence becomes understandable because of the requirement $0.5 < Q^2/p_T^2 < 2$. Although here p_T is measured in the lab system this cut restricts severely the phase space for resolved photons.

With the concept of resolved photons, experimentally observed phenomena in DIS, such as the forward jet cross section and the 2+1 jet rate can be well described, provided a reasonable large scale for the hard scattering process

is used. This concept and its implementation into a full hadron level Monte Carlo generator can be seen as a model alternative to BFKL calculations¹⁷. The next step would be to search for regions in phase space where the two descriptions do not agree but the data can only be described by one of them.

Acknowledgment

I am grateful for many fruitful discussions with L. Jönsson and H. Küster.

Without the great time and fun I still have with Antje it would not have been possible to search for free living pomerinos on the Canary Islands.

References

1. A. Mueller, *Nucl. Phys. B (Proc. Suppl.)* **18C** (1990) 125.
2. A. Mueller, *J. Phys. G* **17** (1991) 1443.
3. H1 Collaboration, C. Adloff et al., Contributed paper, pa 03-49, ICHEP96, Warsaw, Poland, July 1996 (unpublished).
4. E. Mirkes, D. Zeppenfeld, Fixed-Order QCD Background to BFKL Dynamics in Forward Jet Production, in *Proceedings of the Workshop on Future Physics at HERA*, edited by A. De Roeck, G. Ingelman, R. Klanner (1995).
5. H1 Collaboration, T. Ahmed et al., Contributed paper, Abstract 247, HEP97, Jerusalem, Israel, August 1997 (unpublished).
6. L. Lönnblad, *Comp. Phys. Comm.* **71** (1992) 15, ARIADNE 4.08.
7. H. Jung, The RAPGAP Monte Carlo for Deep Inelastic Scattering, version 2.06, <http://www-h1.desy.de/~jung/rapgap.html>.
8. G. Ingelman, A. Edin, J. Rathsman, *Comp. Phys. Comm.* **101** (1997) 108, LEPTO 6.5.
9. H. Jung, *Comp. Phys. Comm.* **86** (1995) 147.
10. M. Glück, E. Reya, M. Stratman, *Phys. Rev. D* **54** (1996) 5515.
11. G. Schuler, T. Sjöstrand, *Phys. Lett. B* **380** (1996) 105.
12. M. Drees, R. Godbole, *Phys. Rev. D* **50** (1994) 3124.
13. J. Chyla, J. Cvach, Virtual Photon Structure from jet production at HERA, in *Proceedings of the Workshop on Future Physics at HERA*, edited by G. Ingelman, A. De Roeck, R. Klanner (Hamburg, 1996).
14. M. Klasen, G. Kramer, B. Potter, hep-ph/9703302.
15. M. Glück, E. Reya, A. Vogt, *Z. Phys. C* **53** (1992) 127.
16. M. Glück, E. Reya, A. Vogt, *Phys. Lett. B* **306** (1993) 391.
17. J. Bartels et al., *Phys. Lett. B* **384** (1996) 300, hep-ph/9604272.